


# EQUINE REGIONAL SCREEN COMPONENTS

## MIDATLANTIC, UNITED STATES



Delaware  
 D. C.  
 Kentucky  
 Maryland  
 North Carolina  
 Tennessee  
 Virginia  
 West Virginia



Fungi	Weeds	Trees	Indoors	Foods
Smut Mix	Ragweed	Cottonwood/Aspen	Pine Shavings	Alfalfa
Curvularia	Pigweed/Careless Weed	Willow	Redwood Shavings	Barley
Cladosporium	Lamb's Quarters	Ash	Cedar Shavings	Corn
Helminthosporium	Cocklebur	Juniper/Cedar	Leather	Cottonseed
Stemphylium	English Plantain	Box Elder/Maple	Sisal	Molasses
Candida Albicans	Dock/Sheep Sorrel	Oak	Wool	Milo
Monilia	Sage	Elm	Cotton	Oats
Cephalosporium	Marsh Elder	Pecan/Hickory		Pasture (Mix)
Mucor Mix	Waterhemp	Walnut		Soybean
Trichoderma		Mulberry	<b>Insects</b>	Wheat
Rhizopus		Sycamore	Cockroach	Rice Bran
Chaetomium	<b>Grasses</b>	Birch	Culicoides	Beet Pulp
Phoma	Blue/June Grass		Mosquito	
Rhodotorula	Orchard	<b>Dust</b>	Stable Fly	<b>Grains</b>
Botrytis	Fescue	Barn Dust	Deer Fly	Corn Pollen
Epicoccum	Bermuda		Hose Fly	Oat Pollen
Fusarium	Redtop	<b>Epidermals</b>	Fire Ant	Wheat Pollen
Pullularia	Timothy	Mouse Epithelium		Barley Pollen
Aspergillus	Ryegrass			
Nigrospora	Bahia			
Alternaria	Johnson Grass			
Penicillium	Vernal			
Saccharomyces				
Cephalothecium				