

ALLERGENIC EXTRACT PRESCRIPTION PRODUCT

FOR ANIMAL HYPOSENSITIZATION
RESTRICTED TO USE BY OR UNDER THE
SUPERVISION OF A VETERINARIAN

INDICATIONS AND USAGE

Keep treatment sets refrigerated at 2-7C. Do not freeze.
Injections are administered subcutaneously using sterile
needle and syringes.

NEVER INJECTION INTRAVENOUSLY

Before injecting, withdraw the syringe plunger slightly. If
blood appears in the syringe, reinsert in another site. The
hyposensitization program may vary with each individual
animal. The following injection schedule serves as a
guideline. The actual dosage should always be determined
by the veterinarian.

HANDLING ADVERSE REACTIONS, IF OBSERVED

Animals should be observed for at least 30 minutes
following an injection. Possible systemic reactions to watch
for are: increased itching, vomiting and/or diarrhea and hives.
The possibility of an anaphylactic reaction is remote. However,
if such a reaction occurs, it may be advisable to administer
epinephrine. For an average size dog, injection 0.2 to 0.5ml of
epinephrine 1:1000 w/v. Mild reactions may be controlled with
antihistamines.

INSTRUCTIONS FOR ADMINISTRATION OF REFILL VIAL(S):

This schedule is to be followed after completion of a three vial
treatment series:

DAY 1	0.5ml (30 days after last injection)
DAY 15	1.0ml

CONTINUE WITH A 1.0ml INJECTION ONCE A MONTH

If the animal is on a two treatment set regimen, give both
injections on the same day at different sites.

In the even of a reaction such as vomiting, diarrhea or increased
itching, call your veterinarian immediately.

Spectrum Labs
2143 E 5th St
Tempe, AZ 85281

**UNITED STATES VETERINARY LICENSE
NUMBER 403**

CALL 800-553-1391